

May 1998

SINP/TNP/98-13
hep-ph/9806214

New constraints on R -parity violation from proton stability

Gautam Bhattacharyya* and Palash B. Pal†

Saha Institute of Nuclear Physics, 1/AF Bidhan Nagar, Calcutta 700064, India

Abstract

We derive stringent upper bounds on all the $(\lambda''_{ijk}\mu_l)$ -type combinations from the consideration of proton stability, where λ''_{ijk} are baryon-number-violating trilinear couplings and μ_l are lepton-number-violating bilinear mass parameters in a R -parity-violating supersymmetric theory.

In the standard model (SM) of particle interactions, baryon and lepton numbers (B and L respectively) appear as accidental global symmetries which are not violated in any order of perturbation theory. On the other hand, in the minimally supersymmetrized version of this model (MSSM), unless one *assumes* that those are indeed conserved quantities, R -parity-violating ($\mathcal{R}$) couplings are naturally allowed [1]. Defined as $R = (-1)^{(3B+L+2S)}$ (where S is the spin of the particle), R -parity is a discrete symmetry under which SM particles are even while their superpartners are odd. In the MSSM, the most general superpotential is expressed as

$$W = W_0 + W', \quad (1)$$

where W_0 and W' represent the R -parity-conserving and R -parity-violating interactions, respectively. In terms of the MSSM superfields, W_0 and W' can be expressed as

$$W_0 = f_e^{ij} L_i H_d E_j^c + f_d^{ij} Q_i H_d D_j^c + f_u^{ij} Q_i H_u U_j^c + \mu H_d H_u, \quad (2)$$

$$W' = \frac{1}{2} \lambda_{ijk} L_i L_j E_k^c + \lambda'_{ijk} L_i Q_j D_k^c + \frac{1}{2} \lambda''_{ijk} U_i^c D_j^c D_k^c + \mu_i L_i H_u, \quad (3)$$

where i, j, k are generation indices, which are assumed to be summed over. In eqs. (2) and (3), L_i and Q_i are SU(2)-doublet lepton and quark superfields, E_i^c, U_i^c, D_i^c are SU(2)-singlet charged lepton, up- and down-quark superfields, and H_d and H_u are Higgs superfields that are responsible for the down- and up-type masses respectively. In eq. (3), λ, λ' and μ_i are L -violating while λ'' are B -violating parameters. λ_{ijk} is antisymmetric under the interchange of the first two family indices, while λ''_{ijk} is antisymmetric under the interchange of the last two. Thus there could be 27 λ' -type, 9 each of λ - and λ'' -type couplings and 3 μ_i parameters when R -parity is explicitly broken.

Since no evidence of L or B violation has been found in experiments to date, one can take either of the two attitudes. The first is to impose R -parity that forbids all the terms in W' . The second is to take the view that since there is no good theoretical motivation for applying such a symmetry *a priori*, perhaps it is more general to admit all interactions invariant under supersymmetry and gauge symmetry, and to examine what are the bounds on the couplings in W' from various phenomenological considerations [2]. We take the latter approach in this paper.

The prime phenomenological concern attached to any $\mathcal{R}$ theory is the question of proton stability. Proton decay requires *both* L and B violations. Therefore the nonobservation of proton decay could be translated to bounds on the simultaneous presence of L - and B -violating terms. Thus what could be obtained in this way are correlated bounds involving the λ'' with one of the λ', λ or μ_i . Investigations along this direction have already revealed the following constraints.

*E-mail address: gb@tnp.saha.ernet.in

†E-mail address: pbpal@tnp.saha.ernet.in




Figure 1: Proton decay mediated by the B -violating λ''_{112} in association with the L -violating μ_l at the tree level. ψ_l is the lepton doublet.

(i) The simultaneous presence of λ' and λ'' couplings involving the lighter generations drives proton decay ($p \rightarrow \pi^0 e^+$) at tree level yielding extremely tight constraints. From the bound on the proton lifetime $\tau_p \gtrsim 10^{32} \text{ y}$ in the given channel, one obtains [3], for an exchanged squark mass of 1 TeV,

$$\lambda'_{11k} \lambda''_{11k} \lesssim 10^{-24}, \quad (4)$$

where $k = 2, 3$. These constraints weaken linearly with the squark mass as the latter increases.

(ii) One can always find at least one diagram at one loop level in which any λ'_{ijk} in conjunction with any λ''_{lmn} contributes to proton decay [4]. It follows that for an exchanged scalar mass of 1 TeV,

$$\lambda'_{ijk} \lambda''_{lmn} \lesssim 10^{-9}. \quad (5)$$

If one admits tree level flavour-changing squark mixing, the bounds are strengthened by two orders of magnitude.

(iii) The combination of λ'' and λ , in association with the charged lepton Yukawa coupling f_e , drives $p \rightarrow K^+ \ell^\pm \ell'^\mp \bar{\nu}$ at tree level [5]. It should be noted though that the analysis in [5] is based on an extended $\text{SU}(3)_c \times \text{SU}(3)_L \times \text{U}(1)$ gauge model. However the argument applies also to the usual $\text{SU}(3)_c \times \text{SU}(2)_L \times \text{U}(1)$ gauge theory, and the same bound holds. In absence of any direct experimental limits on the above decay modes, this bound is

$$\lambda''_{112} \lambda_{ijk} \lesssim 10^{-16} \quad (k \neq 3), \quad (6)$$

assuming a benchmark value $\tau_p \gtrsim 10^{31} \text{ y}$ and an exchanged scalar mass of 1 TeV.

What therefore remains to be done is to examine the bounds on the third and final type of L - and B -violating product couplings $\lambda''_{ijk} \mu_l$. This is precisely our goal in the present paper.

Consider the mechanism of nucleon decay shown in Fig. 1. Here one utilizes the B -violating λ''_{112} , the L -violating μ_l and the canonical R -parity-conserving Yukawa coupling f_u . For the sake of convenience, we assume that all flavour indices correspond to particles in their mass basis. Recalling that λ''_{ijk} is antisymmetric under the interchange of the last two indices, at the quark level we obtain the processes

$$d^c s^c \rightarrow u \bar{\nu}_l, \quad d^c s^c \rightarrow d \bar{l}. \quad (7)$$

They imply the following decay modes of the proton:

$$p \rightarrow K^+ \nu_l, \quad p \rightarrow K^+ \pi^+ l^-, \quad (8)$$

where the charged lepton in the final state can only be e or μ . The relevant dimension-6 operators have an effective coupling

$$G_R \simeq \frac{\lambda''_{112} f_u}{m_{\tilde{u}_R}^2} \left(\frac{\mu_l}{m_{\tilde{H}_u}} \right), \quad (9)$$

leading to an approximate proton lifetime

$$\tau_p^R \simeq (m_p^5 G_R^2)^{-1}. \quad (10)$$




Figure 2: Loop-driven proton decay mediated by the B -violating λ''_{ijk} in association with the L -violating μ_l .

Among the decay modes in eq. (8), the channel $p \rightarrow K^+ \nu$ ($\tau_p \gtrsim 10^{32} y$ [6]) offers the most stringent constraints. For a squark mass $m_{\tilde{u}_R} = 1 \text{ TeV}$, we obtain

$$\lambda''_{112} \epsilon_l \lesssim 10^{-21}, \quad (11)$$

where ϵ_l is defined through $\mu_l \equiv \epsilon_l \mu$. For our order of magnitude estimate, we have made a simple approximation $\mu \simeq m_{\tilde{H}_u} \simeq 1 \text{ TeV}$.

In the mass basis that we have employed, λ''_{112} is the only independent baryon number violating coupling that can be bounded in conjunction with μ_l at the tree level. However, the other λ'' couplings also drive proton decay at loop levels, and therefore are also constrained. This can be seen from the 1-loop diagram of Fig. 2, where h^+ denotes the physical charged Higgs, given by

$$h^+ = \sin \beta H_d^+ - \cos \beta H_u^+, \quad (12)$$

where $\tan \beta \equiv \langle H_u^0 \rangle / \langle H_d^0 \rangle$. Let us consider the case of a neutrino in the final state. In this case, the quark on the line q_k must be a down-type quark. The index k cannot be 3 since there cannot be a b -quark in the final state. For $k = 1$ and 2, the final states will be $\pi^+ \bar{\nu}$ and $K^+ \bar{\nu}$. Indeed, with $k = 1$ and 2, we can cover all the independent λ''_{ijk} -couplings because of the antisymmetry of these couplings in the last two indices. The effective coupling for the dimension-6 operator is given by

$$G'_H \simeq \left(\frac{f_u^i f_d^j}{16\pi^2} \right) (V_{i1}^* V_{1j}) \frac{\lambda''_{ijk} f_d^k}{m_{\tilde{d}_R}^2} \left(\frac{\mu^2}{m_{\tilde{H}}^2} \right) \epsilon_l, \quad (13)$$

where $\tilde{H}$ is a linear combination of $\tilde{H}_u$ and $\tilde{H}_d$. Although this diagram exists for λ''_{112} as well, more stringent bounds on $\lambda''_{112} \epsilon_l$ originate from the tree level diagram discussed earlier. For all other combinations, the bounds are displayed in Table 1.

Before we conclude, a few comments are in order. Although it has been argued [7] that the $L_i H_u$ terms can be rotated away by suitable redefinitions of the L_i and H_d fields in the superpotential, they reappear through quantum corrections after supersymmetry is broken. Moreover, λ - and λ' -couplings generate bilinear terms via one-loop graphs [8]. Phenomenological constraints on μ_l originate from the

Table 1: Summary of upper bounds on the products $\lambda''_{ijk} \epsilon_l$ derived in this paper. The index i appears as row-headings. The indices j, k appear as column headings. The table entry gives the upper bound, which are independent of the index l , for superparticle masses of order 1 TeV.

	21	31	32
1	10^{-21}	10^{-10}	10^{-11}
2	10^{-13}	10^{-12}	10^{-13}
3	10^{-14}	10^{-13}	10^{-14}

fact that the bilinear terms trigger sneutrino-Higgs and consequently neutrino-neutralino mixing, leading to non-zero neutrino masses [9]. Similarly, individual constraints on various λ'' exist. They arise, e.g., from $n-\bar{n}$ oscillation [10] and precision electroweak observables [11].

To conclude, we have used the constraints on proton stability to obtain stringent upper limits on all products of the form $\lambda''_{ijk}\mu_l$. These are complementary to the bounds on the products of the L - and B -violating couplings obtained before. The combinations $\lambda''_{112}\mu_l$ contribute to proton decay at the tree level and are therefore very suppressed. The other combinations drive proton decay through loop diagrams and the resulting constraints are also displayed. Most importantly, the parameter space we have examined in this paper has never been explored before.

References

- [1] G. Farrar and P. Fayet, *Phys. Lett.* **B76** (1978) 575; S. Weinberg, *Phys. Rev.* **D26** (1982) 287; N. Sakai and T. Yanagida, *Nucl. Phys.* **B197** (1982) 533; C. Aulakh and R. Mohapatra, *Phys. Lett.* **B119** (1982) 136; L. Hall and M. Suzuki, *Nucl. Phys.* **B231** (1984) 419; J. Ellis *et al.*, *Phys. Lett.* **B150** (1985) 142; G. Ross and J. Valle, *Phys. Lett.* **B151** (1985) 375; S. Dawson, *Nucl. Phys.* **B261** (1985) 297; R. Barbieri and A. Masiero, *Nucl. Phys.* **B267** (1986) 679.
- [2] For reviews, see G. Bhattacharyya, [hep-ph/9709395](#), *Nucl. Phys. Proc. Suppl.* **52A** (1997) 83; H. Dreiner, [hep-ph/9707435](#).
- [3] I. Hinchliffe and T. Kaeding, *Phys. Rev.* **D47** (1993) 279.
- [4] A.Y. Smirnov and F. Vissani, *Phys. Lett.* **B380** (1996) 317. See also, C. Carlson, P. Roy and M. Sher, *Phys. Lett.* **B357** (1995) 99.
- [5] H. N. Long, P.B. Pal, [hep-ph/9711455](#).
- [6] Review of Particle Physics, *Phys. Rev.* **D54** (1996) 1.
- [7] Hall and Suzuki, Ref. [1].
- [8] S. Roy and B. Mukhopadhyaya, *Phys. Rev.* **D55** (1997) 7020.
- [9] F. de Campos, M. A. Garcia-Jareno, A. S. Joshipura, J. Rosiek, J. W. F. Valle, *Nucl. Phys.* **B451** (1995) 3.
- [10] F. Zwirner, *Phys. Lett.* **B132** (1983) 103; J. L. Goity and M. Sher, *Phys. Lett.* **B346** (1995) 69, Erratum *ibid* **B385** (1996) 500.
- [11] G. Bhattacharyya, D. Choudhury, K. Sridhar, *Phys. Lett.* **B355** (1995) 193.